

- J. Olanzapine-induced weight gain is associated with the -759C/T and -697G/C polymorphisms of the HTR2C gene. *Pharmacogenomics Journal* 2009; 9: 234–241.
61. Gurevich EV, Bordelon Y, Shapiro RM, Arnold SE, Gur RE, Joyce JN. Mesolimbic dopamine D3 receptors and use of antipsychotics in patients with schizophrenia. A postmortem study. *Arch Gen Psychiatry* 1997; 54: 225–232.
62. Reynolds G, Yao Z, Zhang X, Sun J, Zhang Z. Pharmacogenetics of treatment in first-episode schizophrenia: D3 and 5-HT2C receptor polymorphisms separately associate with positive and negative symptom response. *European Neuropsychopharmacology* 2005; 15: 143–151.
63. Adams D, Close S, Farmen M, Downing A, Breier A, Houston J. Dopamine receptor D3 genotype association with greater acute positive symptom remission with olanzapine therapy in predominately caucasian patients with chronic schizophrenia or schizoaffective. *Human Psychopharmacology-Clinical and Experimental* 2008; 23: 267–274.
64. Reynolds GP, Templeman LA, Zhang ZJ. The role of 5-HT2C receptor polymorphisms in the pharmacogenetics of antipsychotic drug treatment. *Progress in Neuro-Psychopharmacology & Biological Psychiatry* 2005; 29: 1021–1028.
65. Samanaite R, Gillespie A, Sendt KV, McQueen G, MacCabe JH, Egerton A. Biological Predictors of Clozapine Response: A Systematic Review. *Front Psychiatry* 2018; 9: 327.
66. Zhang JP, Malhotra AK. Recent Progress in Pharmacogenomics of Antipsychotic Drug Response. *Curr Psychiatry Rep* 2018; 20: 24.
67. Miller CA, Sweatt JD. Covalent modification of DNA regulates memory formation. *Neuron* 2007; 53: 857–869.
68. Miller CA, Gavin CF, White JA, Parrish RR, Honasoge A, Yancey CR, Rivera IM, et al. Cortical DNA methylation maintains remote memory. *Nature Neuroscience* 2010; 13: 664–666.
69. Ghadirivasfi M, Nohešara S, Ahmadkhaniha HR, Eskandari MR, Mostafavi S, Thiagalingam S, Abdolmaleky HM. Hypomethylation of the Serotonin Receptor Type-2 A Gene (HTR2A) at T102C Polymorphic Site in DNA Derived From the Saliva of Patients With Schizophrenia and Bipolar Disorder. *American Journal of Medical Genetics Part B-Neuropsychiatric Genetics* 2011; 156 B: 536–545.
70. Kebir O, Chaumette B, Fatjó-Vilas M, Ambalavanan A, Ramoz N, Xiong L, Mouaffak F, et al. Family-based association study of common variants, rare mutation study and epistatic interaction detection in HDAC genes in schizophrenia. *Schizophrenia Research* 2014; 160: 97–103.
71. Levenson JM, O'Riordan KJ, Brown KD, Trinh MA, Molfese DL, Sweatt JD. Regulation of histone acetylation during memory formation in the hippocampus. *Journal of Biological Chemistry* 2004; 279: 40545–40559.
72. Peleg S, Sananbenesi F, Zovolis A, Burkhardt S, Bahari-Javan S, Agis-Balboa RC, Cota P, et al. Altered Histone Acetylation Is Associated with Age-Dependent Memory Impairment in Mice. *Science* 2010; 328: 753–756.
73. Alarcon JM, Malleret G, Touzani K, Vronskaya S, Ishii S, Kandel ER, Barco A. Chromatin acetylation, memory, and LTP are impaired in CBP^{+/−} mice: A model for the cognitive deficit in Rubinstein-Taybi syndrome and its amelioration. *Neuron* 2004; 42: 947–959.
74. Chuang DM, Leng Y, Marinova Z, Kim HJ, Chiu CT. Multiple roles of HDAC inhibition in neurodegenerative conditions. *Trends in Neurosciences* 2009; 32: 591–601.
75. Weiwer M, Lewis MC, Wagner FF, Holson EB. Therapeutic potential of isoform selective HDAC inhibitors for the treatment of schizophrenia. *Future Med Chem* 2013; 5: 1491–1508.
76. Kebir O, Chaumette B, Fatjó-Vilas M, Ambalavanan A, Ramoz N, Xiong L, Mouaffak F, et al. Family-based association study of common variants, rare mutation study and epistatic interaction detection in HDAC genes in schizophrenia. *Schizophrenia Research* 2014; 160: 97–103.
77. Kim T, Park JK, Kim HJ, Chung JH, Kim JW. Association of histone deacetylase genes with schizophrenia in Korean population. *Psychiatry Res* 2010; 178: 266–269.
78. Brousse G, Meary A, Blanc O, Lancon C, Llorca P, Leboyer M. Clinical predictors of response to olanzapine or risperidone during acute episode of schizophrenia. *Psychiatry Research* 2010; 179: 12–18.
79. Cheah SY, Lawford BR, Young RM, Morris CP, Voisey J. mRNA Expression and DNA Methylation Analysis of Serotonin Receptor 2 A (HTR2A) in the Human Schizophrenic Brain. *Genes (Basel)* 2017; 8.
80. Ghadirivasfi M, Nohešara S, Ahmadkhaniha HR, Eskandari MR, Mostafavi S, Thiagalingam S, Abdolmaleky HM. Hypomethylation of the serotonin receptor type-2 A Gene (HTR2A) at T102C polymorphic site in DNA derived from the saliva of patients with schizophrenia and bipolar disorder. *Am J Med Genet B Neuropsychiatr Genet* 2011; 156 B: 536–545.
81. Dazzan P. Neuroimaging biomarkers to predict treatment response in schizophrenia. *European Neuropsychopharmacology* 2016; 26: S140–S140.
82. Dazzan P. Neuroimaging biomarkers to predict treatment response in schizophrenia: the end of 30 years of solitude? *Dialogues in Clinical Neuroscience* 2014; 16: 491–503.

