

## LITERATURA

- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders (DSM-5). Arlington, VA: American Psychiatric Publishing. 2013.
- World Health Organization. International Classification of Diseases, 11th Revision (ICD-11). Geneva: WHO. 2019.
- Kozinszky Z, Dudas RB. Validation studies of the Edinburgh Postnatal Depression Scale for the antenatal period. *J Affect Disord.* 2015;176:95-105.
- American College of Nurse-Midwives. A Brief Depression Screening Tool for Perinatal Clinical Practice. 2022.
- Yin X, Sun N, Jiang N, et al. Prevalence and associated factors of antenatal depression: systematic reviews and meta-analyses. *Clin Psychol Rev.* 2021;83:101932.
- Boekhorst MGBM, Beerthuis A, Endendijk JJ, et al. Different trajectories of depressive symptoms during pregnancy: a longitudinal population-based study in China. *J Affect Disord.* 2019;248:139-146.
- Bao C, Jin D, Sun S, Xu L, et al. Trajectories and depressive symptoms during the perinatal period: a longitudinal population-based study in China. 2022;13:762719.
- Lee H, Kim KE, Kim MY, et al. Trajectories of depressive symptoms and anxiety during pregnancy and associations with pregnancy stress. *Int J Environ Res Public Health.* 2021;18(5):2733.
- Gelaye B, Sanchez SE, Andrade A, et al. Association of antepartum depression, generalized anxiety, and posttraumatic stress disorder with infant birth weight and gestational age at delivery. *J Affect Disord.* 2020;262:310-316.
- Rahi M, Heikkinen T, Hartter S, et al. Placental transfer of quetiapine in relation to P-glycoprotein activity. *J Psychopharmacol.* 2007;21(7):751-756.
- Ewing G, Tatarchuk Y, Appleby D, et al. Placental transfer of antidepressant medications: implications for postnatal adaptation syndrome. *Clin Pharmacokinet.* 2015;54(4):359-370.
- Watanabe O. Current evaluation of teratogenic and fetotoxic effects of psychotropic drugs. *Seishin Shinkeigaku Zasshi.* 2014;116(12):996-1004.
- Betcher HK, Wisner KL. Psychotropic treatment during pregnancy: research synthesis and clinical care principles. *J Womens Health (Larchmt).* 2020;29(3):310-318.
- Seifertová D, Mohr P, Strunzová V. Léčba psychofarmaky v těhotenství a laktaci. *Psychiatrie Pro Praxi.* 2007;3:118-124.
- Šebela A, Nosková E, Goetz M, et al. Psychofarmaka během těhotenství – mírní, nebo zvyšují riziko pro plod? Část druhá: stabilizátory nálady, anxiolytika. *Pediatric Pro Praxi.* 2017;18(5):282-286.
- Cepeda MS, Kern DM, Nicholson S. Treatment resistant depression in women with peripartum depression. *BMC Pregnancy Childbirth.* 2019;19(1):323.
- Ornoy A, Weinstein-Fudim L, Ergaz Z. Antidepressants, antipsychotics, and mood stabilizers in pregnancy: what do we know and how should we treat pregnant women with depression. *Birth Defects Res.* 2017;109(12):933-956.
- Gentile S. Use of benzodiazepines in pregnancy and lactation: a review. *Expert Opin Drug Saf.* 2011;10(4):537-551.
- Shyken JM, Babbar S, et al. Benzodiazepines in pregnancy. *Clin Obstet Gynecol.* 2019;62(1):156-167.
- Tomson T, Battino D, Bonizzoni E, et al. Comparative risk of major congenital malformations with eight different antiepileptic drugs: a prospective cohort study of the EURAP registry. *Lancet Neurol.* 2018;17(6):530-538.
- Česká psychiatrická společnost. Doporučené postupy pro léčbu depresivních poruch v těhotenství a při kojení. Praha: Česká psychiatrická společnost. 2022.
- Málová V. Esketamin. *Psychiatr. praxi.* 2023;24(3):194-198.
- Meltzer-Brody S, Kanes S. Brexanolone injection in postpartum depression: two multicentre, double-blind, randomised, placebo-controlled, phase 3 trials. *Lancet.* 2018;392(10152):1058-1070.
- Grigoriadis S, VonderPorten EH, Mamisashvili L, et al. The impact of maternal depression during pregnancy on perinatal outcomes: a systematic review and meta-analysis. *J Clin Psychiatry.* 2013;74(4):e321-e341.
- Grote NK, Bridge JA, Gavin AR, et al. A meta-analysis of depression during pregnancy and the risk of preterm birth, low birth weight, and intrauterine growth restriction. *Arch Gen Psychiatry.* 2010;67(10):1012-1024.
- Jarde A, Morais M, Kingston D, et al. Neonatal outcomes in women with untreated antenatal depression compared with women without depression: a systematic review and meta-analysis. *JAMA Psychiatry.* 2016;73(8):826-837.
- Yang J, Qu Y, Zhan Y, et al. Trajectories of depressive symptoms during pregnancy and risk of premature birth: a multicenter and prospective cohort study. *Psychiatry Res.* 2023;326:115251.
- Staneva A, Bogossian F, Pritchard M, et al. The effects of maternal depression, anxiety, and perceived stress during pregnancy on preterm birth: a systematic review. *Midwifery.* 2015;28(3):179-193.
- Ghimire U, Papabathini SS, Kwaku J, et al. Depression during pregnancy and the risk of low birth weight, preterm birth and intrauterine growth restriction: an updated meta-analysis. *Early Hum Dev.* 2021;152:105243.
- Shenassa ED, Widemann LG, Hunt CD. Antepartum depression and preterm birth: pathophysiology, epidemiology, and disparities due to structural racism. *Curr Psychiatry Rep.* 2021;23(3):14.
- Jahan N, Went TR, Sultan W, et al. Untreated depression during pregnancy and its effect on pregnancy outcomes: a systematic review. *Cureus.* 2021;13(8):e17251.
- Shapiro GD, Fraser WD, Frasch MG, et al. Psychosocial stress in pregnancy and preterm birth: associations and mechanisms. *J Perinat Med.* 2013;41(6):631-645.
- Bindt C, Guo N, Bonle MT, et al. Association of antepartum and postpartum depression in Ghanaian and Ivorian women with febrile illness in their offspring: a prospective birth cohort study. *Am J Epidemiol.* 2013;178(9):1394-1402.
- Bonari L, Pinto N, Ahn E, et al. Perinatal risks of untreated depression during pregnancy. *Can J Psychiatry.* 2004;49(11):726-735.
- Mesches GA, Wisner KL, Betcher HK. A common clinical conundrum: antidepressant treatment of depression in pregnant women. *Semin Perinatol.* 2020;44(3):151-229.
- Grigoriadis S, Graves L, Peer M, et al. Maternal anxiety during pregnancy and the association with adverse perinatal outcomes: systematic review and meta-analysis. *J Clin Psychiatry.* 2018;79(5):17r1201.
- Gentile S. Untreated depression during pregnancy: short- and long-term effects in offspring. A systematic review. *Neuroscience.* 2017;342:154-166.
- Weikum WM, Mayes LC, Grunau RE, et al. The impact of prenatal serotonin reuptake inhibitor (SSRI) antidepressant exposure and maternal mood on mother-infant interactions at 3 months of age. *Infant Behav Dev.* 2013;36(4):485-493.
- Rogers A, Obst S, Teague SJ, et al. Association between maternal perinatal depression and anxiety and child and adolescent development: a meta-analysis. *JAMA Pediatr.* 2020;174(11):1082-1092.
- Grigoriadis S, VonderPorten EH, Mamisashvili L, et al. Antidepressant exposure during pregnancy and congenital malformations: is there an association? A systematic review and meta-analysis of the best evidence. *J Clin Psychiatry.* 2013;74(4):e293-e308.
- Bérard A, Zhao JP, Sheehy O. Antidepressant use during pregnancy and the risk of major congenital malformations in a cohort of depressed pregnant women: an updated analysis of the Quebec Pregnancy Cohort. *BMJ Open.* 2017;7(1):e013372.
- Bar-Oz B, Einarson T, Einarson A, et al. Paroxetine and congenital malformations: meta-analysis and consideration of potential confounding factors. *Clin Ther.* 2007;29(5):918-926.
- Yang J, Qu Y, Zhan Y, et al. Trajectories of depressive symptoms during pregnancy and risk of premature birth: a multicenter and prospective cohort study. *J Affect Disord.* 2020;275:149-156.
- Venkatesh KK, Castro VM, Perlis RH, et al. Impact of antidepressant treatment during pregnancy on obstetric outcomes among women previously treated for depression: an observational cohort study. *J Perinatol.* 2017;37(9):1003-1009.
- Malm H, Sourander A, Gissler M, et al. Pregnancy complications following prenatal exposure to SSRIs or maternal psychiatric disorders: results from population-based national register data. *Am J Psychiatry.* 2015;172(12):1224-1231.
- Huang H, Coleman S, Bridge JA, et al. A meta-analysis of the relationship between antidepressant use in pregnancy and the risk of preterm birth and low birth weight. *Hosp Psychiatry.* 2014;36(1):13-18.
- Prady SL, Hanlon I, Fraser LK, et al. A systematic review of maternal antidepressant use in pregnancy and short- and long-term offspring outcomes. *Arch Womens Ment Health.* 2018;21(2):127-140.
- Gurnot C, Martin-Subero I, Mah SM, et al. Prenatal antidepressant exposure associated with CYP2E1 DNA methylation change in neonates. *Epigenetics.* 2015;10(5):361-368.
- El Marroun H, White T, Verhulst FC, et al. Maternal use of antidepressant or anxiolytic medication during pregnancy and childhood neurodevelopmental outcomes: a systematic review. *Eur Child Adolesc Psychiatry.* 2014;23(10):973-992.
- Korayem GB, Alanazi Y, Alanazi H, et al. Exploring women's perception and attitude towards antidepressant use: a cross-sectional study. *BMC Womens Health.* 2024;24(1):453.
- Ban L, Tata LJ, Fiaschi L, et al. Live and non-live pregnancy outcomes among women with depression and anxiety: a population-based study. *PLoS One.* 2012;7(8):e43462.
- Adhikari K, Patten SB, Lee S, et al. Risk of adverse perinatal outcomes among women with pharmacologically treated and untreated depression during pregnancy: a retrospective cohort study. *Paediatr Perinat Epidemiol.* 2019;33(5):323-331.
- Marks C, Silvola R, Teal E, et al. Comparing newborn outcomes after prenatal exposure to individual antidepressants: a retrospective cohort study. *Pharmacotherapy.* 2021;41(11):907-914.
- Tharp MA, Silvola RM, Marks C, et al. Does lack of exposure to individual antidepressants at different points during pregnancy associate with reduced risk of adverse newborn outcomes? *BMC Pregnancy Childbirth.* 2022;22(1):926.
- Fitton CA, Steiner MFC, Aucott L, et al. In utero exposure to antidepressant medication and neonatal and child outcomes: a systematic review. *Acta Psychiatr Scand.* 2020;141(1):21-33.
- Ng QX, Venkatanarayanan N, Ho CYX, et al. Selective serotonin reuptake inhibitors and persistent pulmonary hypertension of the newborn: an updated meta-analysis. *J Womens Health (Larchmt).* 2019;28(3):331-338.
- Huybrechts KF, Bateman BT, Palmsten K, et al. Antidepressant use late in pregnancy and risk of persistent pulmonary hypertension of the newborn. *JAMA.* 2015;313(21):2142-2149.
- Bérard A, Sheehy O, Zhao JP, et al. SSRI and SNRI use during pregnancy and the risk of persistent pulmonary hypertension of the newborn. *Br J Clin Pharmacol.* 2017;83(5):1126-1133.
- Wurst KE, Poole C, Ephross SA, et al. First trimester paroxetine use and the prevalence of congenital, specifically cardiac, defects: a meta-analysis of epidemiological studies. *Birth Defects Res A Clin Mol Teratol.* 2010;88(3):159-170.
- Grigoriadis S, Graves L, Peer M, et al. Benzodiazepine use during pregnancy alone or in combination with an antidepressant and congenital malformations: systematic review and meta-analysis. *J Clin Psychiatry.* 2019;80(4):18r12412.
- Chan JKN, Lee KCK, Wong CSM, et al. Prenatal antidepressant use and risk of congenital malformations: a population-based cohort study. *Psychiatry Res.* 2024;339:116038.
- Campbell KSJ, Collier AC, Irvine MA, et al. Maternal serotonin reuptake inhibitor antidepressants have acute effects on fetal heart rate variability in late gestation. *Front Psychiatry.* 2021;12:680177.